

Kootenai Clinic Endoscopy 1919 Lincoln Way, Suite 416 Coeur d'Alene, ID 83814 Phone 208–625–4601 Fax 208–625–4602 kootenaiclinic.org

SIGMOIDOSCOPY PREP INSTRUCTIONS

Patient:	Gastroenterologist:	
Date of test: _	Time to arrive:	
Your ar	rival time may need to be adjusted based on the	doctor's schedule
Where to check i	n:	
Kootenai Clinic Endoscopy 4th Floor (1919 Lincoln Way, Suite 416)		
Specialty P	rocedures (2003 Kootenai Health Way - North hospit	al entrance registration desk)
• If your test amount o	pressure and seizure medications: is in the morning, take your heart, blood pressure of water <u>at least</u> one hour before your test. is in the afternoon, take your medications at your us	
_ Inhalers: Use as pres	scribed, please bring inhalers with you to your appoir	ntment.
_ Blood thinne Stop	rs: days prior to tes	t.
_ Stop Fish Oil	, Flax Seed Oil, THC Oil and CBD Oil three days բ	prior to test.
Oral diabetic r	medications:	
Do not take the morning dose on the day of your test.		**For low blood sugar - drink regular sugared clear soda (Sprite, 7-UP, etc.)
_ Insulin Instruc	etions:	(i, i, i
	<u>A DRIVER IF SEDATION IS USED</u> e for your test and recovery may vary, we will giv	e you an estimated time that your driver will
	be here.	
	ask your driver to be flexible as to the time to pick you	•
•	river needs to leave, please let us know how we can Rideshares (Uber, Lyft, etc.) are not allowed due to	
Bring reading gla		
Bring photo ID ar	nd insurance cards.	
	PLEASE NOTE: THERE MAY BE SEPARATE CHARGES FOR YOUR TEST; THE DOCTOR, THE FACILITY, ANESTHESIA AND PATHOLOGY	
Scheduled by:		

SEE REVERSE SIDE FOR PREP INSTRUCTIONS



SIGMOIDOSCOPY PREP INSTRUCTIONS

BEFORE YOU BEGIN

- Purchase the following medications over the counter (no prescription needed)
 - TWO sodium phosphate/saline enemas (Fleets brand or generic okay)
 - Do not use Mineral Oil enemas.

DAY OF YOUR TEST

- At Home use 1 **Enema** 2 hours before leaving for your test.
- At Home use 2nd Enema 1 hour before leaving for your test.

IF YOU ARE HAVING SEDATION FOR THIS TEST:

- NO solid food after midnight
- You may drink clear liquids up to 4 hours prior to your test
- DO NOT DRINK ANY LIQUIDS AFTER
- NO Gum, NO Hard Candy, and NO Chewing Tobacco for 8 Hours Prior to Test,
- Nothing RED, BLUE OR PURPLE

CLEAR LIQUID DIET:

- Coffee or tea with any sweetener no creamer
- Clear fruit juices apple, white grape or white cranberry
- Gatorade, other clear bottled drinks
- Any clear carbonated soft drinks (7-UP, Sprite, etc.)
- Clear broth Chicken, beef or vegetable
- Jello: orange, lemon, lime, pineapple or peach (nothing added)
- Water

Please call the Endoscopy Clinic if you have any concerns about this process at (208) 625–4601.

*******Cancellations within 3 business days may be subject to a \$75.00 charge*******